

Propositional Calculus

- *Semantics* (3/3)

Moonzoo Kim
CS Division of EECS Dept.
KAIST

moonzoo@cs.kaist.ac.kr
<http://pswlab.kaist.ac.kr/courses/cs402-07>

Overview

- 2.1 Boolean operators
- 2.2 Propositional formulas
- 2.3 Interpretations
- 2.4 Logical Equivalence and substitution
- 2.5 Satisfiability, validity, and consequence
- 2.6 Semantic tableaux
- 2.7 Soundness and completeness

Semantic tableaux

- The method of **semantic tableaux** is a relatively efficient algorithm for deciding **satisfiability** in the propositional calculus.
 - Search systematically for a model.
 - If one is found, the formula is satisfiable; otherwise, it is unsatisfiable.
- This method is the main tool for proving general theorems about the calculus.

Semantic tableaux

Definition 2.43

- A **literal** is an atom or a negation of an atom.
- An atom is a positive literal and the negation of an atom is a negative literal.
- For any atom p , $\{p, \neg p\}$ is a **complementary** pair of literals.
- For any formula A , $\{A, \neg A\}$ is a **complementary** pair of formulas.
- A is the complement of $\neg A$ and $\neg A$ is the complement of A .

Semantic tableaux

- Analyze the satisfiability of $A = p \wedge (\neg q \vee \neg p)$
 $\nu(A) = T$ iff both $\nu(p) = T$ and $\nu(\neg q \vee \neg p) = T$.
Hence, $\nu(A) = T$ if and only if either:
 1. $\nu(p) = T$ and $\nu(\neg q) = T$ or
 2. $\nu(p) = T$ and $\nu(\neg p) = T$ $\rightarrow \{p, \neg p\}$ or $\{p, \neg q\}$

Reduce the question of the satisfiability of formula A to question about the satisfiability of sets of **literals**.
(top-down approach)

Semantic tableaux



- Formula $B = (p \vee q) \wedge (\neg p \wedge \neg q)$.
- $\nu(B) = T$ iff $\nu(p \vee q) = T$ and $\nu(\neg p \wedge \neg q) = T$.
- Hence, $\nu(B) = T$ iff $\nu(p \vee q) = \nu(\neg p) = \nu(\neg q) = T$.
- Hence, $\nu(B) = T$ iff either
 1. $\nu(p) = \nu(\neg p) = \nu(\neg q) = T$, or
 2. $\nu(q) = \nu(\neg p) = \nu(\neg q) = T$.

Since both $\{p, \neg p, \neg q\}$ and $\{q, \neg p, \neg q\}$ contain **complementary** pairs, B is unsatisfiable.



Semantic tableaux

- The systematic search is easy to conduct if a data structure is used to keep track of the assignments that must be made to subformulas.
- In semantic tableaux, *trees* are used.
- A leaf containing a complementary set of literals will be marked \times , while a satisfiable leaf will be marked \odot .

Semantic tableaux



Semantic tableaux



Semantic tableaux

- α -formulas are conjunctive and are satisfiable only if both subformulas α_1 and α_2 are satisfied
- β -formulas are disjunctive and are satisfied even if only one of the subformulas β_1 or β_2 is satisfiable.

α	α_1	α_2
$\neg\neg A_1$	A_1	
$A_1 \wedge A_2$	A_1	A_2
$\neg(A_1 \vee A_2)$	$\neg A_1$	$\neg A_2$
$\neg(A_1 \rightarrow A_2)$	A_1	$\neg A_2$
$\neg(A_1 \uparrow A_2)$	A_1	A_2
$A_1 \downarrow A_2$	$\neg A_1$	$\neg A_2$
$A_1 \leftrightarrow A_2$	$A_1 \rightarrow A_2$	$A_2 \rightarrow A_1$
$\neg(A_1 \oplus A_2)$	$A_1 \rightarrow A_2$	$A_2 \rightarrow A_1$

β	β_1	β_2
$\neg(B_1 \wedge B_2)$	$\neg B_1$	$\neg B_2$
$B_1 \vee B_2$	B_1	B_2
$B_1 \rightarrow B_2$	$\neg B_1$	B_2
$B_1 \uparrow B_2$	$\neg B_1$	$\neg B_2$
$\neg(B_1 \downarrow B_2)$	B_1	B_2
$\neg(B_1 \leftrightarrow B_2)$	$\neg(B_1 \rightarrow B_2)$	$\neg(B_2 \rightarrow B_1)$
$B_1 \oplus B_2$	$\neg(B_1 \rightarrow B_2)$	$\neg(B_2 \rightarrow B_1)$

Semantic tableaux

- Algorithm 2.46 (Construction of a semantic tableau)

Input: A formula A of the propositional calculus.

Output: A semantic tableau \mathcal{T} for A all of whose leaves are marked.

\mathcal{T} for A is a tree each node of which will be labeled with a set of formulas

$U(l)$: the set of formula of leaf l .

The construction terminates when all leaves are marked \times or \odot .

Semantic tableaux

- If $U(l)$ is a set of literals, check if there is a complementary pair of literals in $U(l)$. If so, mark the leaf closed \times ; if not, mark the leaf as open \odot .
- If $U(l)$ is not a set of literals, **choose a formula** in $U(l)$ which is **not** a literal.
 - If the formula is an α -formula, create a new node l' as a child of l and label l' with $U(l') = (U(l) - \{\alpha\}) \cup \{\alpha_1, \alpha_2\}$.
 - If the formula is a β -formula, create two new nodes l' and l'' as children of l . Label l' with $U(l') = (U(l) - \{\beta\}) \cup \{\beta_1\}$, and label l'' with $U(l'') = (U(l) - \{\beta\}) \cup \{\beta_2\}$.

Semantic tableaux

Definition 2.47

- A tableau whose construction has terminated is called a *completed tableau*.
- A completed tableau is **closed** if all leaves are marked closed (x). Otherwise, it is **open**.


Theorem 2.48

- The construction of a semantic tableau terminates.


Soundness and completeness

- A tool is **sound** if the tool says that a formula ϕ is valid (validity, **not** satisfiability), then ϕ is really valid
- A tool is **complete** if ϕ is valid, then the tool says that ϕ is valid
 - Writing in a counter positive way
 - A tool (or method) is **complete** if the tool says that ϕ is not valid, then ϕ is really not valid
- Therefore, if a tool is sound and complete, then
 - the tool says that ϕ is valid iff ϕ is really valid
- Note that
 - if a dumb tool **always** says that ϕ is not valid, then that tool is still sound
 - if a dumb tool **always** says that ϕ is valid, then that tool is still complete

universe of formulas



universe of formulas



Soundness and completeness

Theorem 2.49(Soundness and completeness)

- Let \mathcal{T} be a completed tableau for a formula A . A is unsatisfiable **if and only if** \mathcal{T} is closed.
- Corollary 2.50 A is satisfiable if and only if \mathcal{T} is open.
- Corollary 2.51 A is valid iff the tableau for $\neg A$ closes.
- Corollary 2.52 The method of semantic tableaux is a decision procedure for validity in the propositional calculus.

Soundness

Proof of soundness

- if the tableau \mathcal{T} for a formula A closes, then A is unsatisfiable.
- if a subtree rooted at node n of \mathcal{T} closes, then the set of formulas $U(n)$ labeling n is unsatisfiable.
 - h : height of the node n in \mathcal{T} .
 - If $h = 0$, n is a leaf. Since \mathcal{T} closes, $U(n)$ contains a complementary set of literals. Hence $U(n)$ is unsatisfiable.

Soundness

- If $h > 0$, then some α - or β - rule was used in creating the child(ren) of n :
 - Case 1: An α -rule was used. $U(n) = \{A_1 \wedge A_2\} \cup U_0$ and $U(n') = \{A_1, A_2\} \cup U_0$ for some set of formulas U_0 . But the height of n' is $h-1$, so $U(n')$ is unsatisfiable since the subtree rooted at n' closes. Let ν be an arbitrary interpretation. Since $U(n')$ is unsatisfiable, $\nu(A') = F$ for some $A' \in U(n')$. There are three possibilities:
 - For some $A_0 \in U_0$, $\nu(A_0) = F$. But $A_0 \in U_0 \subseteq U(n)$.
 - $\nu(A_1) = F$. $\nu(A_1 \wedge A_2) = F$, and $A_1 \wedge A_2 \in U(n)$.
 - $\nu(A_2) = F$. $\nu(A_1 \wedge A_2) = F$, and $A_1 \wedge A_2 \in U(n)$.
 Thus $\nu(A) = F$ for some $A \in U(n)$; $U(n)$ is unsatisfiable.




Soundness

- Case 2:

A β -rule was used. $U(n) = \{B_1 \vee B_2\} \cup U_0$, $U(n') = \{B_1\} \cup U_0$, and $U(n'') = \{B_2\} \cup U_0$. By the inductive hypothesis, both $U(n')$ and $U(n'')$ are unsatisfiable. Let ν be an arbitrary interpretation. There are two possibilities:

- $U(n')$ and $U(n'')$ are unsatisfiable because $\nu(B_0) = F$ for some $B_0 \in U_0$. But $B_0 \in U_0 \subseteq U(n)$.
- Otherwise, $\nu(B_0) = T$ for all $B_0 \in U_0$. Since both $U(n')$ and $U(n'')$ are unsatisfiable, $\nu(B_1) = \nu(B_2) = F$. By definition of ν on \vee , $\nu(B_1 \vee B_2) = F$, and $B_1 \vee B_2 \in U(n)$.

Thus $\nu(B) = F$ for some $B \in U(n)$; since ν was arbitrary, $U(n)$ is unsatisfiable.



Completeness

Proof of completeness

- If A is unsatisfiable then every tableau for A closes.
- Contrapositive statement
 - If some tableau for A is open (i.e., if some tableau for A has an open branch), then the formula A is satisfiable.

Completeness

Definition 2.57

- Let U be a set of formulas. U is a **Hintikka** set iff:
 1. For all atoms p appearing in a formula of U , either $p \in U$ or $\neg p \in U$.
 2. If $\alpha \in U$ is an α -formula, then $\alpha_1 \in U$ and $\alpha_2 \in U$.
 3. If $\beta \in U$ is a β -formula, then $\beta_1 \in U$ or $\beta_2 \in U$.

Theorem 2.59

Let l be an open leaf in a completed tableau \mathcal{T} .

Let $U = \bigcup_i U(i)$, where i runs over the set of nodes on the branch from the root to l . Then U is a Hintikka set.

Completeness

Theorem 2.60(Hintikka's Lemma)

- Let U be a Hintikka set. Then U is satisfiable.

Proof of completeness:

- Let \mathcal{T} be a completed open tableau for A . Then U , the union of the labels of the nodes on an open branch, is a Hintikka set by Theorem 2.59 and a model can be found for U by Theorem 2.60. Since A is the formula labeling the root, $A \in U$, so the interpretation is a model of A .