Examples of First Order Theories

Edited slides from the original slides from CS156 by Prof. Z.Manna

CS156: The Calculus of Computation

Zohar Manna

Autumn 2008

Chapter 3: First-Order Theories

First-Order Theories

	Theory	Quantifiers Decidable	QFF Decidable
T_E	Equality	–	✓
T_{PA}	Peano Arithmetic	–	–
$T_{\mathbb{N}}$	Presburger Arithmetic	✓	✓
$T_{\mathbb{Z}}$	Linear Integer Arithmetic	✓	✓
$T_{\mathbb{R}}$	Real Arithmetic	✓	✓
$T_{\mathbb{Q}}$	Linear Rationals	✓	✓
T_{cons}	Lists	–	✓
T_{cons}^{E}	Lists with Equality	–	✓

Theory of Equality T_E I

Signature:

$$\Sigma_= : \{=, a, b, c, \cdots, f, g, h, \cdots, p, q, r, \cdots\}$$

consists of

- $=$, a binary predicate, interpreted with meaning provided by axioms
- all constant, function, and predicate symbols

Axioms of T_E

1. $\forall x.\ x = x$ (reflexivity)
2. $\forall x, y.\ x = y \ \rightarrow\ y = x$ (symmetry)
3. $\forall x, y, z.\ x = y \wedge y = z \ \rightarrow\ x = z$ (transitivity)
4. for each positive integer n and n-ary function symbol f,
   $\forall x_1, \ldots, x_n, y_1, \ldots, y_n.\ \bigwedge_i x_i = y_i$
   $\rightarrow\ f(x_1, \ldots, x_n) = f(y_1, \ldots, y_n)$ (function congruence)

Theory of Equality T_E II

5. for each positive integer n and n-ary predicate symbol p,
   $\forall x_1, \ldots, x_n, y_1, \ldots, y_n.\ \bigwedge_i x_i = y_i$
   $\rightarrow\ (p(x_1, \ldots, x_n) \leftrightarrow p(y_1, \ldots, y_n))$ (predicate congruence)

(function) and (predicate) are axiom schemata.

Example:

(function) for binary function f for $n = 2$:

$$\forall x_1, x_2, y_1, y_2.\ x_1 = y_1 \wedge x_2 = y_2\ \rightarrow\ f(x_1, x_2) = f(y_1, y_2)$$

(predicate) for unary predicate p for $n = 1$:

$$\forall x, y.\ x = y\ \rightarrow\ (p(x)\ \leftrightarrow\ p(y))$$

Note: we omit "congruence" for brevity.

Decidability of T_E I

T_E is undecidable.

The quantifier-free fragment of T_E is decidable. Very efficient algorithm.

Semantic argument method can be used for T_E

Example: Prove

$$F: \ a = b \wedge b = c \ \rightarrow \ g(f(a), b) = g(f(c), a)$$

is T_E-valid.

Decidability of T_E II

Suppose not; then there exists a T_E-interpretation I such that $I \not\models F$. Then,

1.	I	$\not\models$	F	assumption
2.	I	$\models$	$a = b \wedge b = c$	1, $\rightarrow$
3.	I	$\not\models$	$g(f(a), b) = g(f(c), a)$	1, $\rightarrow$
4.	I	$\models$	$a = b$	2, $\wedge$
5.	I	$\models$	$b = c$	2, $\wedge$
6.	I	$\models$	$a = c$	4, 5, (transitivity)
7.	I	$\models$	$f(a) = f(c)$	6, (function)
8.	I	$\models$	$b = a$	4, (symmetry)
9.	I	$\models$	$g(f(a), b) = g(f(c), a)$	7, 8, (function)
10.	I	$\models$	$\bot$	3, 9 contradictory

F is T_E-valid.

Natural Numbers and Integers

Natural numbers	$\mathbb{N} = \{0, 1, 2, \cdots\}$
Integers	$\mathbb{Z} = \{\cdots, -2, -1, 0, 1, 2, \cdots\}$

Three variations:

- Peano arithmetic T_{PA}: natural numbers with addition, multiplication, $=$
- Presburger arithmetic $T_{\mathbb{N}}$: natural numbers with addition, $=$
- Theory of integers $T_{\mathbb{Z}}$: integers with $+, -, >, =$, multiplication by constants

1. Peano Arithmetic T_{PA} (first-order arithmetic)

$$\Sigma_{\text{PA}} : \{0,\ 1,\ +,\ \cdot,\ =\}$$

Equality Axioms: (reflexivity), (symmetry), (transitivity), (function) for $+$, (function) for $\cdot$.

And the axioms:

1. $\forall x.\ \neg(x + 1 = 0)$ (zero)
2. $\forall x, y.\ x + 1 = y + 1 \ \rightarrow\ x = y$ (successor)
3. $F[0] \wedge (\forall x.\ F[x] \ \rightarrow\ F[x + 1]) \ \rightarrow\ \forall x.\ F[x]$ (induction)
4. $\forall x.\ x + 0 = x$ (plus zero)
5. $\forall x, y.\ x + (y + 1) = (x + y) + 1$ (plus successor)
6. $\forall x.\ x \cdot 0 = 0$ (times zero)
7. $\forall x, y.\ x \cdot (y + 1) = x \cdot y + x$ (times successor)

Line 3 is an axiom schema.

Example: $3x + 5 = 2y$ can be written using Σ_{PA} as

$$x + x + x + 1 + 1 + 1 + 1 + 1 = y + y$$

Note: we have $>$ and $\geq$ since

$3x + 5 > 2y$ write as $\exists z.\ z \neq 0 \wedge 3x + 5 = 2y + z$

$3x + 5 \geq 2y$ write as $\exists z.\ 3x + 5 = 2y + z$

Example:

Existence of pythagorean triples (F is T_{PA}-valid):

$$F : \exists x, y, z.\ x \neq 0 \wedge y \neq 0 \wedge z \neq 0 \wedge x \cdot x + y \cdot y = z \cdot z$$

2. Presburger Arithmetic $T_{\mathbb{N}}$

Signature $\Sigma_{\mathbb{N}}$: $\{0,\ 1,\ +,\ =\}$ no multiplication!

Axioms of $T_{\mathbb{N}}$ (equality axioms, with 1-5):

1. $\forall x.\ \neg(x + 1 = 0)$ (zero)
2. $\forall x, y.\ x + 1 = y + 1 \ \rightarrow\ x = y$ (successor)
3. $F[0] \wedge (\forall x.\ F[x] \ \rightarrow\ F[x + 1]) \ \rightarrow\ \forall x.\ F[x]$ (induction)
4. $\forall x.\ x + 0 = x$ (plus zero)
5. $\forall x, y.\ x + (y + 1) = (x + y) + 1$ (plus successor)

Line 3 is an axiom schema.

$T_{\mathbb{N}}$-satisfiability (and thus $T_{\mathbb{N}}$-validity) is decidable (Presburger, 1929)

3. Theory of Integers $T_{\mathbb{Z}}$

Signature:

$$\Sigma_{\mathbb{Z}}: \{\ldots, -2, -1, 0, 1, 2, \ldots, -3\cdot, -2\cdot, 2\cdot, 3\cdot, \ldots, +, -, >, =\}$$

where

- $\ldots, -2, -1, 0, 1, 2, \ldots$ are constants
- $\ldots, -3\cdot, -2\cdot, 2\cdot, 3\cdot, \ldots$ are unary functions
  (intended meaning: $2 \cdot x$ is $x + x$, $-3 \cdot x$ is $-x - x - x$)
- $+, -, >, =$ have the usual meanings.

Relation between $T_{\mathbb{Z}}$ and $T_{\mathbb{N}}$:

$T_{\mathbb{Z}}$ and $T_{\mathbb{N}}$ have the same expressiveness:

- For every $\Sigma_{\mathbb{Z}}$-formula there is an equisatisfiable $\Sigma_{\mathbb{N}}$-formula.
- For every $\Sigma_{\mathbb{N}}$-formula there is an equisatisfiable $\Sigma_{\mathbb{Z}}$-formula.

$\Sigma_{\mathbb{Z}}$-formula F and $\Sigma_{\mathbb{N}}$-formula G are *equisatisfiable* iff:

$$F \text{ is } T_{\mathbb{Z}}\text{-satisfiable} \quad \text{iff} \quad G \text{ is } T_{\mathbb{N}}\text{-satisfiable}$$

1. Theory of Reals $T_{\mathbb{R}}$

Signature:

$$\Sigma_{\mathbb{R}} : \{0,\ 1,\ +,\ -,\ \cdot,\ =,\ \geq\}$$

with multiplication. Axioms in text.

Example:

$$\forall a, b, c.\ b^2 - 4ac \geq 0 \ \leftrightarrow\ \exists x.\ ax^2 + bx + c = 0$$

is $T_{\mathbb{R}}$-valid.

$T_{\mathbb{R}}$ is decidable (Tarski, 1930)
High time complexity

2. Theory of Rationals $T_{\mathbb{Q}}$

Signature:

$$\Sigma_{\mathbb{Q}}: \{0, 1, +, -, =, \geq\}$$

without multiplication. Axioms in text.

Rational coefficients are simple to express in $T_{\mathbb{Q}}$.

Example: Rewrite

$$\frac{1}{2}x + \frac{2}{3}y \geq 4$$

as the $\Sigma_{\mathbb{Q}}$-formula

$$3x + 4y \geq 24$$

$T_{\mathbb{Q}}$ is decidable Quantifier-free fragment of $T_{\mathbb{Q}}$ is efficiently decidable

Theory of Arrays T_A

Signature:

$$\Sigma_A : \ \{\cdot[\cdot],\ \cdot\langle\cdot \triangleleft \cdot\rangle,\ =\}$$

where

- $a[i]$ binary function –
  read array a at index i ("read(a,i)")
- $a\langle i \triangleleft v\rangle$ ternary function –
  write value v to index i of array a ("write(a,i,v)")

Axioms

1. the axioms of (reflexivity), (symmetry), and (transitivity) of T_E
2. $\forall a, i, j.\ i = j \ \rightarrow\ a[i] = a[j]$ (array congruence)
3. $\forall a, v, i, j.\ i = j \ \rightarrow\ a\langle i \triangleleft v\rangle[j] = v$ (read-over-write 1)
4. $\forall a, v, i, j.\ i \neq j \ \rightarrow\ a\langle i \triangleleft v\rangle[j] = a[j]$ (read-over-write 2)

Note: $=$ is only defined for array elements

$$F: \; a[i] = e \;\rightarrow\; a\langle i \triangleleft e\rangle = a$$

not T_{A}-valid, but

$$F': \; a[i] = e \;\rightarrow\; \forall j.\; a\langle i \triangleleft e\rangle[j] = a[j]\;,$$

is T_{A}-valid.

Also

$$a = b \;\rightarrow\; a[i] = b[i]$$

is not T_{A}-valid: We have only axiomatized a restricted congruence.

T_{A} is undecidable
Quantifier-free fragment of T_{A} is decidable

2. Theory of Arrays $T_{\mathrm{A}}^{=}$ (with extensionality)

Signature and axioms of $T_{\mathrm{A}}^{=}$ are the same as T_{A}, with one additional axiom

$$\forall a, b.\ (\forall i.\ a[i] = b[i]) \ \leftrightarrow\ a = b \quad \text{(extensionality)}$$

Example:

$$F:\ \ a[i] = e \ \rightarrow\ a\langle i \triangleleft e\rangle = a$$

is $T_{\mathrm{A}}^{=}$-valid.

$T_{\mathrm{A}}^{=}$ is undecidable
Quantifier-free fragment of $T_{\mathrm{A}}^{=}$ is decidable